

Khovanov-Rozansky Graph Homology and Composition Product

Emmanuel WAGNER

Abstract

In analogy with a recursive formula for the HOMFLY-PT polynomial of links given by Jaeger, we give a recursive formula for the graph polynomial introduced by Kauffman and Vogel. We show how this formula extends to the Khovanov-Rozansky graph homology.

Introduction

The two variable HOMFLY-PT polynomial P of oriented links in $\mathbb{R}^3$ is uniquely determined by its value on an unknot and by the skein relation in Figure 1, see [3]. The specialization $a = q^n$ and $b = q - q^{-1}$ for a positive integer n gives a Laurent polynomial in one variable q . We denote this one variable polynomial by $P_n(L)$, where L is an oriented link, or by $P_n(D)$ if D is a diagram for L ; the normalization here is

$$P_n(\text{unknot}) = [n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}.$$

Jaeger [5] introduced a recursive formula for the HOMFLY-PT polynomial. In particular, for any oriented link diagram D and any integers $m, n \geq 1$, this formula allows a computation of $P_{n+m}(D)$ as a sum of products $P_n(D_1) P_m(D_2)$ where D_1 and D_2 run over certain subdiagrams of D . Jaeger calls this formula a *composition product*.

In this paper we study finite oriented 4-valent graphs embedded in $\mathbb{R}^2$ such that the orientation of the edges around any vertex is as in Figure 2. We call such graphs in $\mathbb{R}^2$ *regular*. We also allow components of a regular graph to be oriented

$$aP(\text{cross}) - a^{-1}P(\text{cross}) = bP(\text{cup}) P(\text{cap})$$

Figure 1: The HOMFLY-PT skein relation



Figure 2: Local model




Figure 3: Regular graph

circles, see Figure 3 for an example. Expanding all vertices of a regular graph Γ as in Figure 4, Kauffman and Vogel [6] defined for any integer $n \geq 1$ a Laurent polynomial $P_n(\Gamma)$ in one variable q .

In analogy with Jaeger's composition product, we give a formula computing $P_{n+m}(\Gamma)$ as a sum of products $P_n(\Gamma_1) P_m(\Gamma_2)$ where Γ_1 and Γ_2 run over certain regular subgraphs of Γ and $m, n \geq 1$ are integers. More precisely, define a *labelling* of Γ to be a mapping f from the set of edges of Γ to the set $\{1, 2\}$ (an oriented circle is treated as an edge without vertices). We denote $\mathcal{L}(\Gamma)$ the set of labellings of Γ that satisfy the following local condition.

Conservation law: At every vertex v of Γ the number of adjacent edges labelled by 1 (resp. by 2) directed towards v is equal to the number of adjacent edges labelled by 1 (resp. 2) directed out of v .

Given $f \in \mathcal{L}(\Gamma)$, we can erase all edges labelled by 2 (resp. by 1), smooth all 2-valent vertices (see Figure 5) and obtain thus a regular graph denoted $\Gamma_{f,1}$ (resp. $\Gamma_{f,2}$).

Lemma 1. *For all regular graphs $\Gamma \subset \mathbb{R}^2$ and for all integers $m, n \geq 1$,*

$$P_{n+m}(\Gamma) = \sum_{f \in \mathcal{L}(\Gamma)} q^{\sigma(\Gamma, f)} P_n(\Gamma_{f,1}) P_m(\Gamma_{f,2}),$$

where $\sigma(\Gamma, f) = \sigma_{m,n}(\Gamma, f)$ is an integer defined in Section 1.2.

$$\begin{aligned} P_n(\text{X}) &= q P_n(\text{hook}) - q^n P_n(\text{X}) \\ &= q^{-1} P_n(\text{hook}) - q^{-n} P_n(\text{X}) \end{aligned}$$

Figure 4: Graph polynomial P_n



Figure 5: Smoothing of a 2-valent vertex

We consider the categorification of P_n due to Khovanov and Rozansky [8]. Given a regular graph $\Gamma \subset \mathbb{R}^2$ and an integer $n \geq 1$, Khovanov and Rozansky constructed a $\mathbb{Z}$ -graded 2-periodic chain complex $C_n(\Gamma)$ over a commutative polynomial $\mathbb{Q}$ -algebra:

$$C_n(\Gamma) = \bigoplus_{i \in \mathbb{Z}, j \in \mathbb{Z}/2\mathbb{Z}} C_n^{i,j}(\Gamma).$$

The differential d of this complex respects the $\mathbb{Z}$ -grading and increases the $\mathbb{Z}/2\mathbb{Z}$ -grading by one:

$$C_n^{i,j}(\Gamma) \xrightarrow{d} C_n^{i,j+1}(\Gamma).$$

For all $i \in \mathbb{Z}$, $j \in \mathbb{Z}/2\mathbb{Z}$,

$$KR_n^{i,j}(\Gamma) = \text{Ker}(d : C_n^{i,j}(\Gamma) \rightarrow C_n^{i,j+1}(\Gamma)) / \text{Im}(d : C_n^{i,j-1}(\Gamma) \rightarrow C_n^{i,j}(\Gamma))$$

is a finite dimensional vector space over $\mathbb{Q}$. Set for all $i \in \mathbb{Z}$,

$$KR_n^i(\Gamma) = KR_n^{i,0}(\Gamma) \oplus KR_n^{i,1}(\Gamma) \text{ and } KR_n(\Gamma) = \bigoplus_{i \in \mathbb{Z}, j \in \mathbb{Z}/2\mathbb{Z}} KR_n^{i,j}(\Gamma).$$

This construction categorifies the graph polynomial $P_n(\Gamma)$ in the sense that

$$P_n(\Gamma) = \sum_{i \in \mathbb{Z}} \dim_{\mathbb{Q}} KR_n^i(\Gamma) q^i. \quad (1)$$

We will denote by curly brackets $\{.\}$ the shift up of the $\mathbb{Z}$ -grading: for $i, k \in \mathbb{Z}$ and $j \in \mathbb{Z}/2\mathbb{Z}$, $KR_n^{i,j}(\Gamma)\{k\} = KR_n^{i+k,j}(\Gamma)$. For $k \in \mathbb{Z}$, denote $\langle k \rangle$ the shift of the $(\mathbb{Z}/2\mathbb{Z})$ -grading by $k \pmod{2}$. We state now our main result.

Theorem 1. *For all regular graphs $\Gamma \subset \mathbb{R}^2$, for all $m, n \geq 1$, $i \in \mathbb{Z}$ and $j \in \mathbb{Z}/2\mathbb{Z}$,*

$$KR_{n+m}^{i,j}(\Gamma) \cong \bigoplus_{\substack{f \in \mathcal{L}(\Gamma) \\ k, l \in \mathbb{Z}, k+l+\sigma(\Gamma, f) = i \\ r, s \in \mathbb{Z}/2\mathbb{Z}, r+s = j}} KR_n^{k,r}(\Gamma_{f,1}) \otimes_{\mathbb{Q}} KR_m^{l,s}(\Gamma_{f,2}) \{\sigma(\Gamma, f)\}$$

where $\cong$ is a $\mathbb{Q}$ -linear isomorphism and $\sigma(\Gamma, f) = \sigma_{m,n}(\Gamma, f)$ is an integer defined in Section 1.2.

This theorem yields a categorification of Lemma 1 and gives a computation of $KR_{n+m}(\Gamma)$ via $KR_n(\Gamma_{f,1})$ and $KR_m(\Gamma_{f,2})$. We derive from Theorem 1 a direct computational formula for $KR_n(\Gamma)$, see Corollary 1, Section 3.

The plan of the paper is as follows. In the first section we explain all notations of Lemma 1 and Theorem 1. The second section is devoted to the proofs of Lemma 1 and Theorem 1. In the third section, we explore consequences of Theorem 1.

$$P_n(\text{circle with arrow}) = \frac{q^n - q^{-n}}{q - q^{-1}} = [n]_q \quad (2)$$

$$P_n(\text{circle with arrow and a crossing}) = [n-1]_q P_n(\text{circle with arrow}) \quad (3)$$

$$P_n\left(\begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array}\right) = [2]_q P_n(\text{cross}) \quad (4)$$

$$P_n(\text{circle with arrow and a crossing}) = P_n(\text{cup}) + [n-2]_q P_n(\text{circle with arrow}) \quad (5)$$

$$P_n\left(\begin{array}{c} \nearrow \searrow \nearrow \searrow \\ \nwarrow \nearrow \nwarrow \nearrow \end{array}\right) + P_n\left(\begin{array}{c} \nearrow \searrow \nearrow \searrow \\ \nwarrow \nearrow \nwarrow \nearrow \end{array}\right) = P_n\left(\begin{array}{c} \nearrow \searrow \nearrow \searrow \\ \nwarrow \nearrow \nwarrow \nearrow \end{array}\right) + P_n\left(\begin{array}{c} \nearrow \searrow \nearrow \searrow \\ \nwarrow \nearrow \nwarrow \nearrow \end{array}\right) \quad (6)$$

Figure 6: Graph relations

1 Preliminaries

1.1 Graph polynomials

The graph polynomial $P_n(\Gamma) \in \mathbb{Z}[q, q^{-1}]$ ($n \geq 1$) of a regular graph $\Gamma \subset \mathbb{R}^2$ is defined from the relations in Figure 4. Notice that $P_n(\Gamma)$ is preserved under ambient isotopy of Γ in $\mathbb{R}^2$. The polynomial P_n can also be defined as the only polynomial invariant of regular graphs invariant under ambient isotopy of graphs in $\mathbb{R}^2$ and satisfying the relations in Figure 6, see [6]. In other words, these relations are sufficient to compute $P_n(\Gamma)$ recursively. Murakami, Ohtsuki and Yamada [10] gave a state sum formula for $P_n(\Gamma)$ and deduced that $P_n(\Gamma)$ has only non-negative coefficients for any regular graph Γ and any $n \geq 1$.

1.2 Notations

Let $\Gamma \subset \mathbb{R}^2$ be a regular graph. We define the *rotation number* of Γ . Smooth all the vertices of Γ as in Figure 7. This gives a disjoint union of oriented circles embedded in $\mathbb{R}^2$; we call these circles *Seifert circles* of Γ . The rotation number of Γ , denoted $r(\Gamma)$, is the sum of the signs of these circles where the sign of a Seifert circle is +1 if it is oriented counterclockwise and -1 otherwise.

We define the interaction $\langle v|\Gamma|f \rangle \in \mathbb{Z}$ of a vertex v of Γ with a labelling f as




Figure 7: Smoothing




Figure 8: Definition of the interaction $\langle v|\Gamma|f\rangle$

shown in Figure 8, where 1 and 2 are the values of f on the corresponding edges. Set $\langle \Gamma|f\rangle = \sum_v \langle v|\Gamma|f\rangle$, where v runs over all vertices of Γ . Given integers $m, n \geq 1$, set

$$\sigma(\Gamma, f) = \sigma_{m,n}(\Gamma, f) = \langle \Gamma, f\rangle + m r(\Gamma_{f,1}) - n r(\Gamma_{f,2}) \in \mathbb{Z}.$$

2 Proofs

2.1 Proof of Lemma 1

Fix $m, n \geq 1$. For any regular graph $\Gamma \subset \mathbb{R}^2$, set

$$Q(\Gamma) = Q_{n+m}(\Gamma) = \sum_{f \in \mathcal{L}(\Gamma)} q^{\sigma(\Gamma, f)} P_n(\Gamma_{f,1}) P_m(\Gamma_{f,2}) \in \mathbb{Z}[q, q^{-1}].$$

In order to prove the lemma it is enough to check that Q satisfies the five relations on Figure 9. First we verify (7):

$$\begin{aligned} Q(\bigcirc) &= q^{-m} P_n(\bigcirc) + q^n P_m(\bigcirc) \\ &= q^{-m} [n]_q + q^n [m]_q = [n+m]_q. \end{aligned}$$

We need to fix more notations: given a regular graph Γ , a labelling $f_0 \in \mathcal{L}(\Gamma)$, and a subset E_0 of the set of edges of Γ , set

$$Q(\Gamma_{f_0, E_0}) = \sum_{f \in \mathcal{L}(\Gamma), f|_{E_0} = f_0|_{E_0}} q^{\sigma(\Gamma, f)} P_n(\Gamma_{f,1}) P_m(\Gamma_{f,2}).$$

$$Q(\bigcirc) = \frac{q^{n+m} - q^{-(n+m)}}{q - q^{-1}} = [n+m]_q \quad (7)$$

$$Q(\text{graph with one crossing}) = [n+m-1]_q Q(\text{graph with no crossing}) \quad (8)$$

$$Q\left(\begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array}\right) = [2]_q Q(\text{graph with two crossings}) \quad (9)$$

$$Q(\text{graph with two crossings}) = Q(\text{graph with one crossing}) + [n+m-2]_q Q(\text{graph with no crossing}) \quad (10)$$

$$Q\left(\begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array}\right) + Q\left(\begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array}\right) = Q\left(\begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array}\right) + Q\left(\begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array}\right) \quad (11)$$

Figure 9: Graph relations

For example, $Q(\text{graph with one crossing})$ is given by a sum over all labellings whose values on the pictured edges are 1. We now check (8). We have

$$Q(\text{graph with one crossing}) = Q(\text{graph with one crossing}) + Q(\text{graph with one crossing}) + Q(\text{graph with one crossing}) + Q(\text{graph with one crossing}). \quad (12)$$

Applying relation (2) to the P_m -terms in $Q(\text{graph with one crossing})$, we obtain

$$Q(\text{graph with one crossing}) = q^{n-1}[m]_q Q(\text{graph with one crossing}),$$

and applying relation (3) to the P_n -terms in $Q(\text{graph with one crossing})$, we get

$$Q(\text{graph with one crossing}) = q^{-m}[n-1]_q Q(\text{graph with one crossing}).$$

Similarly, we can apply relation (3) to the P_m -terms in $Q(\text{graph with one crossing})$ and relation (2) to the P_n -terms in $Q(\text{graph with one crossing})$ and we get that the right hand-side of (12) is equal to

$$\begin{aligned}
& q^{-m}[n-1]_q Q(\text{ }^1_1) + q^{n-1}[m]_q Q(\text{ }^1_1) \\
& + q^n[m-1]_q Q(\text{ }^2_2) + q^{-m+1}[n]_q Q(\text{ }^2_2) \\
& = (q^{-m}[n-1]_q + q^{n-1}[m]_q) Q(\text{ }^1_1) + (q^n[m-1]_q + q^{-m+1}[n]_q) Q(\text{ }^2_2) \\
& = [n+m-1]_q (Q(\text{ }^1_1) + Q(\text{ }^2_2)) \\
& = [n+m-1]_q Q(\text{ }^1_1).
\end{aligned}$$

Hence Q satisfies (8).

We check (9):

$$\begin{aligned}
Q\left(\text{ }^1_1\right) &= Q\left(\text{ }^1_1\right) + Q\left(\text{ }^2_2\right) + Q\left(\text{ }^1_1\right) \\
&+ Q\left(\text{ }^2_2\right) + Q\left(\text{ }^1_1\right) + Q\left(\text{ }^2_2\right) \\
&+ Q\left(\text{ }^1_1\right) + Q\left(\text{ }^2_2\right) + Q\left(\text{ }^1_1\right) \\
&+ Q\left(\text{ }^2_2\right) \\
&= Q\left(\text{ }^1_1\right) + Q\left(\text{ }^2_2\right) + q^{-1}Q\left(\text{ }^1_1\right) + qQ\left(\text{ }^2_2\right) \\
&+ qQ\left(\text{ }^1_1\right) + q^{-1}Q\left(\text{ }^2_2\right) + qQ\left(\text{ }^1_1\right) \\
&+ q^{-1}Q\left(\text{ }^2_2\right) + q^{-1}Q\left(\text{ }^1_1\right) + qQ\left(\text{ }^2_2\right).
\end{aligned}$$

Using the relation (4) for the P_n -terms and P_m -terms, we easily obtain (9).

We now check (10).

$$\begin{aligned}
Q\left(\text{ }^1_1\right) &= Q\left(\text{ }^1_1\right) + Q\left(\text{ }^2_2\right) \\
&+ Q\left(\text{ }^2_2\right) + Q\left(\text{ }^1_1\right) \\
&+ Q\left(\text{ }^1_1\right) + Q\left(\text{ }^2_2\right) \\
&+ Q\left(\text{ }^2_2\right) + Q\left(\text{ }^1_1\right) \\
&+ Q\left(\text{ }^1_1\right) + Q\left(\text{ }^2_2\right).
\end{aligned}$$

We apply the relations (2), (3), and (5) to the P_n -terms and P_m -terms, so the latter expression is equal to:

$$\begin{aligned}
& Q \left(\begin{array}{c} \curvearrowright_1^1 \\ \end{array} \right) + q^{-m}[n-2]_q Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) + q^{n-2}[m]_q Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) \\
& + Q \left(\begin{array}{c} \curvearrowright_2^2 \\ \end{array} \right) + q^n[m-2]_q Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) + q^{2-m}[n]_q Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) \\
& + q^{1-m}[n-1]_q Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) + q^{n-1}[m-1]_q Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) \\
& + q^{1-n}[m-1]_q Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) + q^{m-1}[n-1]_q Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) \\
& + Q \left(\begin{array}{c} \curvearrowright_1^1 \\ \end{array} \right) + Q \left(\begin{array}{c} \curvearrowright_2^2 \\ \end{array} \right) \\
& = Q \left(\begin{array}{c} \curvearrowright \\ \end{array} \right) + [n+m-2]_q Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right).
\end{aligned}$$

In order to prove the last relation (11), we put in correspondence the labellings occuring on the two sides of (11) as in the following two examples:

$$Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) = Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right), \quad Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) = Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right).$$

There are 28 different labellings possible for . Among these labellings, 14 are identified with labellings of  as in the first example and 8 with labellings of  as in the second example. The remaining 6 labellings are involved in the following equalities (three other equalities are obtained by exchanging 1 and 2):

$$\begin{aligned}
& Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) + Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) = Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) + Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right), \\
& Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) + Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) = Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right), \\
& Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) = Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right) + Q \left(\begin{array}{c} \text{ } \text{ } \text{ } \\ \text{ } \text{ } \text{ } \end{array} \right).
\end{aligned}$$

In this way we obtain that Q satisfies (11). Hence, Q satisfies equations (7-11) and since these equations determine P_{n+m} , we conclude that $Q = P_{n+m}$. Lemma 1 is proved.

2.2 Proof of Theorem 1

Since the graph polynomial has only non negative coefficients, we have directly from Lemma 1 and Formula (1):

$$KR_{n+m}^i(\Gamma) \cong \bigoplus_{\substack{f \in \mathcal{L}(\Gamma) \\ k, l \in \mathbb{Z}, k+l+\sigma(\Gamma, f) = i}} KR_n^k(\Gamma_{f,1}) \otimes_{\mathbb{Q}} KR_m^l(\Gamma_{f,2}) \{\sigma(\Gamma, f)\}. \quad (13)$$

Khovanov and Rozansky [8], p. 78 proved the following: for any regular graph Γ , $n \geq 1$, and $i \in \mathbb{Z}$,

$$KR_n^{i,j}(\Gamma) = 0 \text{ if } j = r(\Gamma) + 1 \pmod{2},$$

where $r(\Gamma)$ is the rotation number defined in Section 1.2. Furthermore, given a regular graph Γ and $f \in \mathcal{L}(\Gamma)$, we have

$$r(\Gamma) = r(\Gamma_{f,1}) + r(\Gamma_{f,2}).$$

Theorem 1 follows from (13) and the latter formula.

3 Consequences

Given a regular graph Γ , we denote $L(\Gamma)$ the subset of labellings of Γ such that $\Gamma_{f,2}$ is a disjoint union of circles. We define also

$$\mathcal{S}(\Gamma) = \{\Gamma_{f,1} | f \in L(\Gamma)\}.$$

Given a regular subgraph $\Delta \in \mathcal{S}(\Gamma)$ of Γ , there is a unique labelling $f_{\Delta} \in L(\Gamma)$ such that $\Delta = \Gamma_{f_{\Delta},1}$. For $\Delta \in \mathcal{S}(\Gamma)$, set $\beta(\Gamma, \Delta) = \langle \Gamma, f_{\Delta} \rangle$ where $\langle \Gamma, f_{\Delta} \rangle$ is defined in Section 1.2. We need to fix more notations. For all $k, l \in \mathbb{Z}$,

$$\mathbb{Q}\{k\}\langle l \rangle = \bigoplus_{i \in \mathbb{Z}, j \in \mathbb{Z}/2\mathbb{Z}} \mathbb{Q}\{k\}\langle l \rangle^{i,j},$$

where

$$\mathbb{Q}\{k\}\langle l \rangle^{i,j} = \begin{cases} \mathbb{Q} & \text{if } i = k \text{ and } j = l \pmod{2}, \\ 0 & \text{otherwise.} \end{cases}$$

We state a corollary of Theorem 1.

Corollary 1. For all regular graphs $\Gamma \subset \mathbb{R}^2$ and all integers $n \geq 2$,

$$KR_n(\Gamma) \cong \bigoplus_{\Delta_1 \in \mathcal{S}(\Gamma), \Delta_2 \in \mathcal{S}(\Delta_1), \dots, \Delta_{n-1} \in \mathcal{S}(\Delta_{n-2})} \mathbb{Q}\{\delta(\Delta_1, \dots, \Delta_{n-1})\} \langle r(\Gamma) \rangle$$

where

$$\begin{aligned} \delta(\Delta_1, \dots, \Delta_{n-1}) &= \sum_{i=0}^{n-2} (\beta(\Delta_i, \Delta_{i+1}) + (n-i) r(\Delta_{i+1}) - (n-1-i) r(\Delta_i)) \\ &= \sum_{i=0}^{n-2} (\beta(\Delta_i, \Delta_{i+1}) + 2 r(\Delta_{i+1})) - (n-1) r(\Gamma) \end{aligned}$$

with the convention $\Delta_0 = \Gamma$.

Proof. From [8] we have

$$KR_1(\Gamma) = \begin{cases} \mathbb{Q} & \text{if } \Gamma \text{ is a union of circles,} \\ 0 & \text{otherwise.} \end{cases}$$

The rest of the argument is just a straightforward recurrence using the case $m = 1$ of Theorem 1. $\square$

We illustrate Theorem 1 and Corollary 1, for $n = 2$ and $m = 1$ in the following example.

Example 1.

$$\begin{aligned} KR_3 \left(\bigcirc \otimes \bigcirc \right) &\cong KR_2 \left(\bigcirc \otimes \bigcirc \right) \oplus KR_2(\bigcirc) \{-3\} \langle 1 \rangle \\ &\quad \oplus KR_2(\bigcirc) \{3\} \langle 1 \rangle \oplus KR_2(\bigcirc) \{1\} \langle 1 \rangle \\ &\quad \oplus KR_2(\bigcirc) \{-1\} \langle 1 \rangle \\ &\cong \mathbb{Q}\{-2\} \oplus \mathbb{Q}\{2\} \oplus \mathbb{Q} \oplus \mathbb{Q} \oplus \mathbb{Q}\{-2\} \oplus \mathbb{Q}\{-4\} \\ &\quad \oplus \mathbb{Q}\{4\} \oplus \mathbb{Q}\{2\} \oplus \mathbb{Q}\{2\} \oplus \mathbb{Q} \oplus \mathbb{Q} \oplus \mathbb{Q}\{-2\} \end{aligned}$$

Remark. Corollary 1 suggest an equivalent but direct definition of $KR_n(\Gamma)$.

$$KR_n(\Gamma) = \bigoplus_{\Delta_1 \in \mathcal{S}(\Gamma), \Delta_2 \in \mathcal{S}(\Delta_1), \dots, \Delta_n \in \mathcal{S}(\Delta_{n-1})} \mathbb{Q}\{\delta(\Delta_1, \dots, \Delta_n)\} \langle r(\Gamma) \rangle.$$

Using this expression as a starting definition of $KR_n(\Gamma)$, it would be interesting to exhibit explicitly the isomorphism of Theorem 1.

Acknowledgments This paper is done as part of the author's PhD research at the Université Louis Pasteur (Strasbourg). The author wishes to thank Prof. Turaev for his endless support.

References

- [1] D. Bar-Natan, *On Khovanov's categorification of the Jones polynomial*, Algebraic and Geometric Topology, 2 (2002), 337-370.
- [2] D. Bar-Natan, *Khovanov's homology for tangles and cobordisms*, math.GT/0410495.
- [3] P. Freyd, D. Yetter, J. Hoste, W. B. R. Lickorish, K. Millet and A. Ocneanu, *A new polynomial invariant of knots and links*, Bull. AMS (N.S.) 12 (1985), 239-246.
- [4] B. Gornik, *Note on Khovanov link cohomology* math.QA/0402266.
- [5] F. Jaeger, *Composition product and models for the Homfly polynomial*, Enseign. Math., 35 (1989), 323-361.
- [6] L. Kauffman and P. Vogel, *Link polynomials and a graphical calculus*, J. Knot Theory Ramifications, 1 (1992), 59-104.
- [7] M. Khovanov, *A categorification of the Jones polynomial*, Duke Math J., 101 (2000), 359-426.
- [8] M. Khovanov and L. Rozansky, *Matrix factorizations and link homology*. math.QA/0401268.
- [9] M. Khovanov and L. Rozansky, *Matrix factorizations and link homology II*. math.QA/0505056.
- [10] H. Murakami, T. Ohtsuki, and S. Yamada, *Homfly polynomial via an invariant of colored plane graphs*. Enseign. Math., 44 (1998), 325-360.
- [11] J. Rasmussen. *Khovanov-Rozansky homology of two-bridge knots and links*. math.GT/0508510.

Institut de Recherche Mathématique Avancée, C.N.R.S.-Université Louis Pasteur, 7 rue René Descartes 67084 Strasbourg Cedex, France.
E-mail address: wagner@math.u-strasbg.fr